

Northumbria Research Link

Citation: Gibbon, Karl, Debusse, Dorothée and Caplan, Nick (2013) Low impact weight-bearing exercise in an upright posture achieves greater lumbopelvic stability than overground walking. *Journal of Bodywork and Movement Therapies*, 17 (4). pp. 462-468. ISSN 1360-8592

Published by: Elsevier

URL: <http://dx.doi.org/10.1016/j.jbmt.2013.02.004>
<<http://dx.doi.org/10.1016/j.jbmt.2013.02.004>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/11150/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

1 **LOW IMPACT WEIGHT-BEARING EXERCISE IN AN UPRIGHT POSTURE**
2 **ACHIEVES GREATER LUMBOPELVIC STABILITY THAN OVERGROUND**
3 **WALKING**
4

5
6
7 ¹Gibbon, K.C., ¹Debusse, D. & ¹Caplan, N.
8

9
10
11 1 Faculty of Health & Life Sciences, Northumbria University, Newcastle upon
12 Tyne, United Kingdom
13

14
15 Corresponding author:
16 Dr Nick Caplan
17 Faculty of Health and Life Sciences
18 Northumbria University
19 Northumberland Building
20 Newcastle upon Tyne
21 NE1 8ST
22 United Kingdom
23

24 Tel: +44 191 243 7382
25 Email: nick.caplan@northumbria.ac.uk
26
27
28

ABSTRACT

The aim of this study was to determine the kinematic differences between movements on a new exercise device (EX) that promotes a stable trunk over a moving, unstable base of support, and overground walking (OW). Sixteen male participants performed EX and OW trials while their movements were tracked using a 3D motion capture system. Trunk and pelvis range of motion (ROM) were similar between EX and OW in the sagittal and frontal planes, and reduced for EX in the transverse plane. The pelvis was tilted anteriorly, on average, by about 16 degrees in EX compared to OW. Hip and knee ROM were reduced in EX compared to OW. The exercise device appears to promote similar or reduced lumbopelvic motion, compared to walking, which could contribute to more tonic activity of the local lumbopelvic musculature.

Keywords: kinematics, walking, lumbopelvic stability, exercise

INTRODUCTION

In vitro studies have shown the thoracolumbar and lumbar spine, devoid of any musculature, will experience structural failure under compressive loadings as small as 20 and 90 N in magnitude, respectively (Crisco et al 1992). Considering spinal loadings experienced *in vivo* can range from 6 kN during selected everyday tasks (McGill & Norman 1986) to in excess of 36 kN during competitive powerlifting (Cholewicki et al 1991) the human vertebral column is intrinsically incapable of meeting the physiological demands placed upon it without additional stabilisation at a segmental level (Panjabi et al 1989).

The role of the lumbopelvic trunk musculature in providing the required supplementary stability at a segmental level is well documented (Bergmark 1989; Panjabi 1992; Cholewicki & McGill 1996; Vera-Garcia et al 2007). In particular, due to their anatomical positioning, morphology and function, the deeper fibres of the lumbar multifidus (LM) and the transversus abdominis (TrA) are considered crucial for local stability of the lumbar spine (Hodges & Richardson 1996; Hodges 1999; Kim et al 2007).

A growing body of evidence links structural and functional changes of local stabilising trunk muscles with low back pain (LBP) (Hides et al 1994; Hides et al 1996; Hodges & Richardson 1996; Danneels et al 2000; Oddsson & De Luca 2003; Hides et al 2008; Hides et al 2008; MacDonald et al 2009; Teyhen et al 2009; Wallwork et al 2009). In people with LBP, muscle fibre atrophy and fatty infiltrations

of the LM have been observed (Kader et al 2000), as well as a dysfunction of the anticipatory activity of the LM and TrA (Hodges & Richardson 1998).

Corrective/restorative treatment strategies for such dysfunction of the local lumbopelvic musculature have included specific motor control exercises (Hides et al 2008), 'core stability' training, muscular strength and endurance training (Danneels et al 2001), aerobic exercise (Frost et al 1995) and the use of an unstable base of support (BOS) (Marshall & Murphy 2006), often in a tailored combination (Demoulin et al 2010). The majority of these approaches tend to show only modest effectiveness (Keller et al 2007; van Middelkoop et al 2010), possibly due to a lack of carry-over to functional day-to-day activities (Richardson & Hides 2004; Hodges & Cholewicki 2007).

Recently a new method promoting activation of LM and TrA has been proposed as an alternative to the current approaches for addressing local lumbopelvic muscle dysfunction (Debuse et al 2013). The users of the exercise device move their feet in a quasi-elliptical path in anti-phase against virtually no external resistance. The absence of external resistance creates the need for much greater motor control of the legs and pelvis, to control leg movement, whilst maintaining an upright trunk posture, than in conventional exercise devices. The exercise device was found to recruit LM and TrA to a greater extent than a range of control activities, including standing on the ground or on an unstable base of support and voluntary muscle contractions. The authors postulated that the method promotes a relatively stable lumbopelvic area during a functional lower limb movement and results in an automatic recruitment/activity of TrA and LM (Debuse et al 2013). Richardson and

Jull in their seminal paper of 1995 proposed that local muscles work tonically, as opposed to global muscles which tend to work phasically. This is widely accepted by other authors working in this field (for example Sahrman 2002; Hides 2004; Hides et al 2004; Hodges & Cholewicki 2007). Debusse et al (2013) imply that tonic muscle activity is likely to be responsible for the stable lumbopelvic region when using the exercise device. However, no information was provided on the lumbopelvic and lower limb kinematics of the user while exercising to identify how the exercise device promoted lumbopelvic stability and, thus, tonic muscle activity.

The aim of the current study was to compare lower limb, pelvic and trunk kinematics during the use of a newly developed exercise device (EX) and overground walking (OW), with a particular focus on the level of lumbopelvic stability in both activities.

METHOD

Participants

Sixteen healthy adult male volunteers (mean \pm SD age: 26.5 \pm 3.38 years, body mass: 82.158 \pm 7.21 kg, height: 1.78 \pm 0.05 m, and body mass index: 25.89 \pm 2.16 kg·m⁻²) with no recent history of LBP, gait impairments, or other conditions affecting their ability to walk or exercise, agreed to participate in this study. Participants gave their fully informed written consent to take part. The study had received ethical approval from the Institutional Review Board prior to data collection.

Three-dimensional Motion Capture

Three-dimensional trajectories of 39 retro-reflective markers ($\varnothing=14\text{mm}$) were captured at a sampling frequency of 200 Hz using a 12 camera near-infrared motion capture facility (MX T20, Vicon Motion Systems, Oxford, UK). Markers were placed in accordance with a standard full-body model (Plug-in-Gait, Vicon Motion Systems, Oxford, UK), which consists of a 15 segment rigid-linked model of the head, thorax, pelvis, and bilateral upper arms, forearms, hands, thighs, lower legs and feet. Only the segmental orientations of the thorax, pelvis, thighs and lower legs were subsequently used for analysis.

The motion capture system was calibrated before all testing sessions using a standard dynamic protocol, with a 5 marker calibration wand (Vicon Motion Systems, Oxford, UK). System calibration was accepted when the image error of all 12 cameras was less than 0.2 mm.

Body mass, height and anthropometric measurements, including leg length (anterior superior iliac spine to medial malleolus), ankle widths and knee widths, necessary for the correct operation of the model used were taken in triplicate and the mean value used thereafter.

Experimental protocol

Participants completed an overground walking (OW) condition and a condition using the exercise device (EX – Figure 1) in a counterbalanced random order within a single session. In the OW condition participants were asked to walk along a level 7.5 m walkway, instrumented with embedded force plates (OR6-7, AMTI, Watertown,

Massachusetts, USA), at a self-selected comfortable speed. Starting positions were adjusted individually to ensure that 'clean' foot contacts with the force plates could be achieved without direct targeting by the participant. A minimum of 10 trials were completed, before six trials - without evidence of targeting - were selected for subsequent analysis.

In the EX condition participants were given an initial five minute period to familiarise themselves with the exercise device. Following this, 30 seconds of trajectory data were captured during exercise in standing. Subsequently, six cycles were chosen at random for analysis. All participants were given standardised instructions on the correct use of the device emphasising the need for a 'slow controlled movement' whilst maintaining 'an upright posture' during each cycle.

Data processing and reduction

Marker trajectories collected during OW and EX trials were reconstructed and processed within Vicon Nexus (1.7, Vicon Motion Systems, Oxford, UK). Lost or obscured trajectory segments were interpolated using a quintic-spline function for gaps less than or equal to 10 frames (0.05 s) or a pattern fill function for gaps less than 10 frames, which utilises the trajectory of a marker with a similar predicted displacement trajectory. Marker trajectories were then low pass filtered at 5 Hz using a fourth-order zero lag Butterworth filter (Saunders et al 2005).

Key "gait cycle" phases (stance and swing) were demarcated for both the OW and EX conditions using discrete gait cycle events. Heel strikes and toe offs during OW were detected using the vertical component of the ground reaction force obtained

from the force plates embedded flush with the walkway surface at the centre of the calibrated capture volume. When using the new exercise device, the feet remain in contact with the foot plates at all times during both stance and swing phase. Therefore, data collected during EX were divided into a stance and swing phase based on the trajectory of a marker placed on the front corner of the foot plate: stance was defined as the most anterior to the most posterior foot plate position, and swing was from the most posterior to most anterior foot plate position.

Three-dimensional angular displacements for the trunk (thorax with respect to [wrt] pelvis), pelvis (wrt the room, rather than a relative position between body segments), hip (pelvis wrt thigh) and knee (thigh wrt lower leg) were time normalised to cycle duration in 2% increments (51 data points from 0-100%) for the right sided cycles of both OW and EX conditions. Angular range of motion (ROM) was calculated as the maximum minus the minimum joint angle achieved within one cycle. This was done for each of the six trials and averaged within each participant, and then between all participants in both conditions. The mean angular position of each segment or joint was determined as the average of each angle throughout the gait cycle for OW and EX. The difference in mean angular positions, or offset, between OW and EX was calculated. Data for each variable were checked for normality of distribution using Q-Q and box plots. For variables that were normally distributed, paired samples t-tests were used to compare ROM and mean angular position between conditions with significance set at $p < 0.05$. For variables that were not normally distributed, Wilcoxon signed rank tests were instead used. Confidence intervals (95%) were also calculated for each pairwise comparison. All statistical analyses were performed using SPSS (version 19).

RESULTS

Spatiotemporal characteristics

All spatiotemporal data were normally distributed. Statistically significant differences were observed in all six spatiotemporal parameters (Table 1). The EX condition was characterised by reduction in cadence ($t=21.220$, $df=15$, $p<0.001$), stride length ($t=14.041$, $df=15$, $p<0.001$), stride duration ($t=26.380$, $df=15$, $p<0.001$), speed ($t=20.506$, $df=15$, $p<0.001$), and effective stance phase ($t=15.354$, $df=15$, $p<0.001$) compared to those observed during OW. Step width was significantly greater in the EX condition compared to OW ($t=2.662$, $df=15$, $p<0.05$).

Kinematics

All angular ROM data were normally distributed with the exception of the hip in the transverse plane. Angular ROM was found to be similar between EX and OW conditions for the trunk in the sagittal ($t=1.622$, $df=15$, $p=0.126$) and frontal ($t=1.203$, $df=15$, $p=0.248$) planes, and was similar for the pelvis in the sagittal ($t=1.607$, $df=15$, $p=0.129$) and frontal ($t=0.213$, $df=15$, $p=0.834$) planes. In the transverse plane, ROM was significantly reduced for the trunk ($t=8.513$, $df=15$, $p<0.001$) and the difference approached significance in the pelvis ($t=1.854$, $df=15$, $p=0.083$) between EX and OW (Table 2).

All mean angular position data were normally distributed with the exception of the pelvis and hip in the transverse plane. The pelvis was significantly tilted anteriorly for the EX condition compared to OW with an offset of 6.49° ($t=4.697$, $df=15$, $p<0.001$) (Table 3). Hip ROM was significantly reduced in the EX condition compared to OW in the sagittal ($t=7.359$, $df=15$, $p<0.001$), frontal ($t=2.572$, $df=15$, $p=0.021$) and transverse ($Z=3.516$, $p<0.001$) planes (Table 2). Knee ROM was also reduced in EX in the sagittal ($t=8.463$, $df=15$, $p<0.001$), frontal ($t=7.041$, $df=15$, $p<0.001$) and transverse ($t=7.120$, $df=15$, $p<0.001$) planes. The hip ($t=13.297$, $df=15$, $p<0.001$) and knee ($t=19.878$, $df=15$, $p<0.001$) were both more flexed throughout the gait cycle in the EX condition than in OW, with offsets of 22.31° and 24.11° , respectively, which were significant (Table 3). Despite the reduced ROM, peak knee and hip angles occurred at a similar point in the gait cycle for OW and EX (Figure 2).

DISCUSSION

The aim of this investigation was to compare the kinematics of lower limb and trunk motion during the use of a newly developed exercise device (EX), and overground walking (OW). The key findings of this study were that the lumbopelvic region was at least as stable whilst exercising on the new exercise device as overground walking. In the transverse plane, reduced ROM was observed during EX compared to OW. This stable lumbopelvic region was achieved over a dynamically moving base of support, where the ROM of the knees and hips was lower in EX than in OW. All

spatiotemporal variables were significantly reduced in EX compared to OW, suggesting a slower, more controlled motion.

Trunk motion in the sagittal and frontal planes demonstrated similar ranges for both EX and OW. In the transverse plane, a reduced ROM was observed for EX suggesting increased lumbopelvic stability. Similar observations were made for the pelvis in terms of ROM, although in the transverse plane, a smaller reduction in range of motion was found for EX, with this reduction approaching statistical significance.

As a fundamental human activity, walking has previously been investigated as an intervention strategy in the treatment of LBP (Torstensen et al 1998; Joffe et al 2002; Taylor et al 2003; Mirovsky et al 2006). However, heterogeneity of study design and methodological quality have contributed to inconsistent findings (Hendrick et al 2010). Of these studies only Torstensen et al. (1998) and Taylor et al. (2003) used walking independently, while Joffe et al. (2002) and Mirovsky et al. (2006) combined walking with bodyweight support and traction, respectively. Notwithstanding the lack of evidence supporting walking as an effective intervention strategy for low back pain, the movement itself, involving control of trunk and pelvis motion during lower limb movements, is known to contribute to recruitment of the TrA and LM (Saunders et al 2004; Saunders et al 2005). Importantly, walking tends to be advocated by health care professionals in line with recommendations that ordinary physical activities should be continued as much as possible in order to aid recovery from LBP and prevent long-term disability (van Tulder et al 2000).

Similarities observed in both trunk and pelvic ROM between EX and OW in the sagittal and frontal planes suggest that the exercise device may be similar to walking, in terms of enabling tonic recruitment of the local lumbopelvic muscles such as TrA and LM. Previously Saunders et al. (2004; 2005) reported tonic TrA but phasic LM activity at walking speeds comparable to those reported here. However, no data were presented describing changes in activity amplitude, if any, within each gait cycle. The phasic activity of LM previously reported during walking (Saunders et al 2004) could be a factor leading to the questionable effectiveness of walking as a successful intervention for LBP (Hendrick et al 2010). The reduced transverse ROM, and thus the inherently more tonic muscle actions, in EX compared with OW seen in the current study could further indicate facilitation of greater tonic activity of the local lumbopelvic muscles (Richardson & Jull 1995) when using the new exercise device than in overground walking. If this reduced axial rotation results in more tonic recruitment of LM at a segmental level, then this could lead to the exercise device being a more successful intervention for LBP than walking. Current research within our group is exploring differences in lumbopelvic muscle recruitment between the exercise device and walking using ultrasound imaging and electromyography. Future studies in symptomatic populations are required to examine the clinical effectiveness of the exercise device.

No angular offsets were found between EX and OW for the trunk or pelvic position in all three planes, with the exception of a greater degree of anterior tilt of the pelvis in the EX condition. Influences of anterior pelvic tilt (O'Sullivan et al 2006) and accompanying lordotic spinal posture (Claus et al 2009), similar in magnitude to that observed within this investigation, have previously been shown to recruit both the

superficial and deep fibres of the LM to approximately 30-40% of maximal voluntary isometric contraction capabilities, which is known to facilitate stabiliser muscle recruitment (McArdle et al 1991). Thus, this angular offset could be beneficial for the recruitment of the LM, provided care is taken to avoid over-recruitment of the superficial fibres of LM.

Hip and knee joints were more flexed throughout the gait cycle in EX than during OW. The increase in hip flexion was partly due to the angular definition being relative to a perpendicular axis of the pelvis. Therefore, the observed increase in anterior tilt creates a greater degree of flexion at the hip. The increased flexion of the knee throughout the gait cycle during EX are linked to the reduced stride length that was caused by the mechanical constraints of the device. By reducing stride length, the knee was unable to reach full extension during the stance phase of the gait cycle, as is normally seen during OW. What was apparent for knee and hip motion in the sagittal plane, was that the change in angle throughout the gait cycle showed a more sinusoidal pattern in EX compared to OW. This, more regular, movement pattern could contribute, to some extent, to more continuous/tonic muscle recruitment, a key training requirement of the local stabilising musculature (Richardson & Jull 1995).

There has been a drive, in recent years, for training interventions for the local muscles of the lumbopelvic region to be made more functional (Hodges 2011). A number of studies have brought into question the transferability of any training effects seen following less functional activities such as gym ball training where the base of support is simply unstable (Drake et al 2006). Debuse et al. (2013) demonstrated that the local lumbopelvic muscles were recruited to a greater extent

with lower limb movement and an unstable base of support than with standing still on an unstable base of support (i.e. no voluntary lower limb movement). While overground walking involves lower limb movement, it does not usually involve an unstable base of support. During exercising on the new device, the requirement to control the descent of the “front” leg by gradually unloading the “back” leg within each gait cycle may result in greater recruitment of the local lumbopelvic muscles than overground walking. Our ultrasound imaging studies will examine this aspect in greater detail.

This study has a number of limitations. It examined relative motion between the pelvis and trunk. In order to gain a better understanding of how the exercise device influences the kinematics of the lumbopelvic region, a more detailed model of the thoracic and lumbar spine is needed. This would enable vertebral motion to be evaluated at a segmental level. Participants were asked to walk at their preferred walking speed. Due to the nature of the exercise device, movements were slower compared to walking. Saunders et al (2005) reported reduced axial rotation of the spine when walking slower. Thus, slow walking could lead to similar kinematics that were observed for the exercise device, and this should be explored further. However, walking slower does not involve an unstable base of support, or the complex motor control associated with using the exercise device, both of which could be contributing to increased local lumbopelvic muscle recruitment.

CONCLUSION

Key differences between exercising on the device and overground walking included reduced transverse plane range of trunk motion with respect to the pelvis (i.e. increased lumbopelvic stability), a more anteriorly tilted pelvis, and reduced stride length, knee and hip range of motion in the sagittal plane. The greater anterior tilt of the pelvis potentially moved the pelvis into a more advantageous position for the recruitment of TrA and LM. However, the unstable base of support afforded by the new exercise device would seem to add a challenge to movement control that may result in greater TrA and LM activity than overground walking. Future investigations should examine TrA and LM activity during walking and exercising on the new device using ultrasound imaging.

REFERENCES

- Bergmark A 1989 Stability of the lumbar spine. A study in mechanical engineering. Acta Orthopaedica Scandinavica. Supplementum 230: 1-54.
- Cholewicki J, McGill SM 1996 Mechanical stability of the in vivo lumbar spine: implications for injury and chronic low back pain. Clinical Biomechanics 11: 1-15.
- Cholewicki J, McGill SM, Norman RW 1991 Lumbar spine loads during the lifting of extremely heavy weights. Medicine and Science in Sports and Exercise 23: 1179-86.
- Claus AP, Hides JA, Moseley GL, Hodges PW 2009 Different ways to balance the spine: subtle changes in sagittal spinal curves affect regional muscle activity. Spine 34: E208-14.
- Crisco JJ, Panjabi MM, Yamamoto I, Oxland TR 1992 Euler Stability of the Human Ligamentous Lumbar Spine Part II: Experiment. Clinical Biomechanics 7: 27-32.
- Danneels LA, Cools AM, Vanderstraeten GG, Cambier DC, Witvrouw EE, Bourgois J, de Cuyper HJ 2001 The effects of three different training modalities on the cross-sectional area of the paravertebral muscles. Scandinavian Journal of Medicine and Science in Sports 11: 335-41.
- Danneels LA, Vanderstraeten GG, Cambier DC, Witvrouw EE, De Cuyper HJ 2000 CT imaging of trunk muscles in chronic low back pain patients and healthy control subjects. European Spine Journal 9: 266-72.
- Debuse D, Birch O, St Clair Gibson A, Caplan N 2013 Low impact weight-bearing exercise in an upright posture increases the activation of two key local

378 muscles of the pumbo-pelvic region. *Physiotherapy Theory and Practice* 29:
379 51-60.

380 Demoulin C, Grosdent S, Capron L, Tomasella M, Somville PR, Crielaard JM,
381 Vanderthommen M 2010 Effectiveness of a semi-intensive multidisciplinary
382 outpatient rehabilitation program in chronic low back pain. *Joint, Bone, Spine :*
383 *Revue du Rhumatisme* 77: 58-63.

384 Drake JDM, Fischer SL, Brown SHM, Callaghan JP 2006 Do Exercise Balls Provide
385 a Training Advantage for Trunk Extensor Exercises? A Biomechanical
386 Evaluation. *Journal of Manipulative and Physiological Therapeutics* 29: 354-
387 362.

388 Frost H, Klaber Moffett JA, Moser JS, Fairbank JC 1995 Randomised controlled trial
389 for evaluation of fitness programme for patients with chronic low back pain.
390 *British Medical Journal* 310: 151-4.

391 Hendrick P, Te Wake A, Tikkisetty A, Wulff L, Yap C, Milosavljevic S 2010 The
392 effectiveness of walking as an intervention for low back pain: a systematic
393 review. *European Spine Journal* 19: 1613-1620.

394 Hides J 2004. Paraspinal mechanism in low back pain. In (Richardson C, Hodges P,
395 Hides J). *Therapeutic exercise for lumbopelvic stabilization*. Churchill
396 Livingstone, Edinburgh. 149-162.

397 Hides J, Gilmore C, Stanton W, Bohlscheid E 2008 Multifidus size and symmetry
398 among chronic LBP and healthy asymptomatic subjects. *Manual Therapy* 13:
399 43-9.

400 Hides J, Richardson C, Hodges P 2004. Local segmental control. In (Richardson C,
401 Hodges P, Hides J). *Therapeutic Exercise for Lumbopelvic Stabilization*.
402 Churchill Livingstone, Edinburgh. 185-220.

403 Hides J, Stanton W, Freke M, Wilson S, McMahon S, Richardson C 2008 MRI study
404 of the size, symmetry and function of the trunk muscles among elite cricketers
405 with and without low back pain. *British Journal of Sports Medicine* 42: 809-13.

406 Hides JA, Richardson CA, Jull GA 1996 Multifidus muscle recovery is not automatic
407 after resolution of acute, first-episode low back pain. *Spine* 21: 2763-9.

408 Hides JA, Stanton WR, McMahon S, Sims K, Richardson CA 2008 Effect of
409 stabilization training on multifidus muscle cross-sectional area among young
410 elite cricketers with low back pain. *Journal of Orthopaedic and Sports Physical
411 Therapy* 38: 101-8.

412 Hides JA, Stokes MJ, Saide M, Jull GA, Cooper DH 1994 Evidence of lumbar
413 multifidus muscle wasting ipsilateral to symptoms in patients with
414 acute/subacute low back pain. *Spine* 19: 165-72.

415 Hodges P, Cholewicki J 2007. Functional control of the spine. In (Vleeming A,
416 Mooney V, Stoeckart R). *Movement, Stability and Lumbopelvic Pain*. Churchill
417 Livingstone, Endinburgh.

418 Hodges PW 1999 Is there a role for transversus abdominis in lumbo-pelvic stability?
419 *Manual Therapy* 4: 74-86.

420 Hodges PW 2011 Pain and motor control: From the laboratory to rehabilitation.
421 *Journal of Electromyography and Kinesiology* 21: 220-8.

422 Hodges PW, Richardson CA 1996 Inefficient muscular stabilization of the lumbar
423 spine associated with low back pain. A motor control evaluation of transversus
424 abdominis. *Spine* 21: 2640-50.

425 Hodges PW, Richardson CA 1998 Delayed postural contraction of transversus
426 abdominis in low back pain associated with movement of the lower limb.
427 *Journal of Spinal Disorders* 11: 46-56.

428 Joffe D, Watkins M, Steiner L, Pfeifer BA 2002 Treadmill ambulation with partial body
429 weight support for the treatment of low back and leg pain. Journal of
430 Orthopaedic and Sports Physical Therapy 32: 202-13.

431 Kader DF, Wardlaw D, Smith FW 2000 Correlation between the MRI changes in the
432 lumbar multifidus muscles and leg pain. Clinical Radiology 55: 145-9.

433 Keller A, Hayden J, Bombardier C, van Tulder M 2007 Effect sizes of non-surgical
434 treatments of non-specific low-back pain. European Spine Journal 16: 1776-
435 88.

436 Kim C, Gottschalk L, Eng C, Ward S, Lieber R 2007 The multifidus muscle is the
437 strongest stabilizer of the lumbar spine. The Spine Journal 7: 76S.

438 MacDonald D, Moseley GL, Hodges PW 2009 Why do some patients keep hurting
439 their back? Evidence of ongoing back muscle dysfunction during remission
440 from recurrent back pain. Pain 142: 183-8.

441 Marshall PW, Murphy BA 2006 Evaluation of functional and neuromuscular changes
442 after exercise rehabilitation for low back pain using a Swiss ball: a pilot study.
443 Journal of Manipulative and Physiological Therapeutics 29: 550-60.

444 McArdle WD, Katch FI, Katch VL 1991. Exercise physiology : energy, nutrition, and
445 human performance. Lea & Febiger, Philadelphia.

446 McGill SM, Norman RW 1986 Partitioning of the L4-L5 dynamic moment into disc,
447 ligamentous, and muscular components during lifting. Spine 11: 666-78.

448 Mirovsky Y, Grober A, Blankstein A, Stabholz L 2006 The effect of ambulatory
449 lumbar traction combined with treadmill on patients with chronic low back
450 pain. Journal of Back and Musculoskeletal Rehabilitation 19: 73-78.

451 O'Sullivan PB, Dankaerts W, Burnett AF, Farrell GT, Jefford E, Naylor CS, O'Sullivan
452 KJ 2006 Effect of different upright sitting postures on spinal-pelvic curvature
453 and trunk muscle activation in a pain-free population. *Spine* 31: E707-12.

454 Oddsson LI, De Luca CJ 2003 Activation imbalances in lumbar spine muscles in the
455 presence of chronic low back pain. *Journal of Applied Physiology* 94: 1410-
456 20.

457 Panjabi M, Abumi K, Duranceau J, Oxland T 1989 Spinal stability and intersegmental
458 muscle forces. A biomechanical model. *Spine* 14: 194-200.

459 Panjabi MM 1992 The stabilizing system of the spine. Part I. Function, dysfunction,
460 adaptation, and enhancement. *Journal of Spinal Disorders* 5: 383-9.

461 Richardson C, Hides J 2004. Closed chain segmental control In (Richardson C,
462 Hodges P, Hides J). *Therapeutic Exercise for Lumbopelvic Stabilization*.
463 Churchill Livingstone, Edinburgh.

464 Richardson CA, Jull GA 1995 Muscle control-pain control. What exercises would you
465 prescribe? *Manual Therapy* 1: 2-10.

466 Sahrmann S 2002. *Diagnosis and Treatment of Movement Impairment Syndromes*.
467 Mosby, St Louis.

468 Saunders SW, Rath D, Hodges PW 2004 Postural and respiratory activation of the
469 trunk muscles changes with mode and speed of locomotion. *Gait and Posture*
470 20: 280-90.

471 Saunders SW, Schache A, Rath D, Hodges PW 2005 Changes in three dimensional
472 lumbo-pelvic kinematics and trunk muscle activity with speed and mode of
473 locomotion. *Clinical Biomechanics* 20: 784-93.

474 Taylor NF, Evans OM, Goldie PA 2003 The effect of walking faster on people with
475 acute low back pain. *European Spine Journal* 12: 166-72.

Teyhen DS, Williamson JN, Carlson NH, Suttles ST, O'Laughlin SJ, Whittaker JL, Goffar SL, Childs JD 2009 Ultrasound characteristics of the deep abdominal muscles during the active straight leg raise test. Archives of Physical Medicine and Rehabilitation 90: 761-7.

Torstensen TA, Ljunggren AE, Meen HD, Odland E, Mowinckel P, Geijerstam S 1998 Efficiency and costs of medical exercise therapy, conventional physiotherapy, and self-exercise in patients with chronic low back pain. A pragmatic, randomized, single-blinded, controlled trial with 1-year follow-up. Spine 23: 2616-24.

van Middelkoop M, Rubinstein SM, Verhagen AP, Ostelo RW, Koes BW, van Tulder MW 2010 Exercise therapy for chronic nonspecific low-back pain. Best Practice and Research. Clinical Rheumatology 24: 193-204.

van Tulder M, Malmivaara A, Esmail R, Koes B 2000 Exercise therapy for low back pain: a systematic review within the framework of the cochrane collaboration back review group. Spine 25: 2784-96.

Vera-Garcia FJ, Elvira JL, Brown SH, McGill SM 2007 Effects of abdominal stabilization maneuvers on the control of spine motion and stability against sudden trunk perturbations. Journal of Electromyography Kinesiology 17: 556-67.

Wallwork TL, Stanton WR, Freke M, Hides JA 2009 The effect of chronic low back pain on size and contraction of the lumbar multifidus muscle. Manual Therapy 14: 496-500.

500 Table 1. Spatiotemporal characteristics of overground walking and exercise in the
 501 standing position on the device. (SD = standard deviation, CI = confidence interval).

502

Gait Parameter	Overground Walking		Exercise Device		Mean Difference	
	Mean	±1SD	Mean	±1SD	(95% CI)	<i>P</i> value
Cadence						
(steps·min ⁻¹)	110.7	7.2	71.3	2.7	-39.4 (-43.4 to -35.45)	<0.001
Stride Length (m)	1.41	0.09	1.10	0.00	-0.31 (-0.35 to -0.26)	<0.001
Stride Duration (s)	1.09	0.07	1.69	0.06	0.60 (0.55 to 0.65)	<0.001
Speed (m·s ⁻¹)	1.30	0.13	0.65	0.03	-0.65 (-0.71 to -0.58)	<0.001
Step Width (m)	0.20	0.03	0.23	0.05	0.03 (0.01 to 0.06)	0.018
Stance Phase (%)	59.54	1.66	49.45	2.26	-10.09 (-11.49 to -8.69)	<0.001

Table 2. Angular range of motion of the trunk, pelvis, hip, and knee in all three planes during overground walking and using the exercise device, also including the mean difference between the two conditions. (SD = standard deviation, CI = confidence interval).

Gait Parameter	Walking		Exercise Device		Mean Difference	
	Mean	±1SD	Mean	±1SD	(95% CI)	<i>P</i> value
Sagittal Plane						
Trunk	3.93	1.80	3.01	1.67	-0.92 (-0.29 to 2.14)	0.126
Pelvis	2.89	0.78	3.69	1.91	0.8 (-1.86 to 0.26)	0.129
Hip	42.54	3.96	33.38	2.28	-9.16 (6.50 to 11.81)	<0.001
Knee	59.88	4.03	45.22	6.02	-14.66 (10.97 to 18.36)	<0.001
Frontal Plane						
Trunk	12.59	3.26	11.21	4.42	-1.39 (-1.07 to 3.84)	0.248
Pelvis	8.29	3.33	8.09	2.70	-0.20 (-1.85 to 2.26)	0.834
Hip	12.67	3.44	8.77	4.64	-3.90 (0.67 to 7.14)	0.021
Knee	16.50	5.91	9.42	5.22	-7.08 (4.93 to 9.22)	<0.001
Transverse Plane						
Trunk	12.55	3.85	3.92	1.14	-8.63 (6.47 to 10.79)	<0.001
Pelvis	12.00	3.28	9.25	4.18	-2.75 (-0.41 to 5.92)	0.083
Hip	16.93	7.34	8.87	2.73	-8.06 (4.95 to 11.17)	<0.001 ^a
Knee	20.66	5.37	10.59	3.96	10.07 (7.06 to 13.09)	<0.001

^a indicates that these data were not normally distributed

Table 3. Mean angular position of the trunk, pelvis, hip and knee in all three planes during overground walking and exercise in the standing position on the device. (SD = standard deviation, CI = confidence interval).

Gait Parameter	Walking		Exercise Device		Mean Difference	
	Mean	±1SD	Mean	±1SD	(±95% CI)	<i>P</i> value
Sagittal Plane						
Trunk	-5.37	6.15	-5.43	6.66	0.06 (-3.44 to 3.56)	0.970
Pelvis	9.06	4.06	15.55	6.18	-6.49 (-9.43 to -3.54)	<0.001
Hip	18.30	5.56	40.61	6.62	-22.31(-25.88 to -18.73)	<0.001
Knee	26.28	4.62	50.39	6.69	-24.11 (-26.69 to -21.52)	<0.001
Frontal Plane						
Trunk	-0.41	1.68	0.53	2.05	-0.94 (-2.27 to 0.38)	0.150
Pelvis	-0.25	1.23	-0.33	1.83	0.08 (-0.69 to 0.85)	0.827
Hip	-0.14	2.02	-0.91	2.33	0.77 (-0.13 to 1.67)	0.088
Knee	2.99	3.92	0.44	7.53	2.55 (-0.37 to 5.48)	0.082
Transverse Plane						
Trunk	-2.11	1.86	-1.70	2.03	-0.41 (-1.25 to 0.42)	0.311
Pelvis	-0.65	2.26	-1.63	2.93	0.98 (-0.05 to 2.01)	0.056 ^a
Hip	8.77	8.35	2.55	5.59	6.22 (1.77 to 10.66)	0.010 ^a
Knee	-8.77	9.14	1.16	8.58	-9.94 (-12.45 to -7.42)	<0.001

^a indicates that these data were not normally distributed

FIGURE CAPTIONS

Figure 1. The exercise device during use.

Figure 2. Hip (A) and knee (B) angles are presented for overground walking (- - -) and exercise (—) conditions. The shaded region represents the standard deviation for the exercise device data series.

Figure 1



530 **Figure 2**
531

